

Beta-Carotene 20% DC



Chemical names of active ingredient

Beta-carotene, provitamin A

CAS-No. 7235-40-7

EINECS-No. 230-636-6

PRD-No.
30061292

Article

50094891 25 kg bag in box

Country of origin
Denmark

Description

Free-flowing, dark-red powder with a mild characteristic odor, consisting of spherical particles with a uniform particle size. Some white starch particles may be visible.

Composition

Ingredients in descending order of weight: gelatin, beta-carotene, corn starch, sucrose, sodium ascorbate, tricalcium phosphate, tocopherol (E306).


Solubility

Dispersible in warm water (35 – 40 °C), to form a stable orange dispersion.

Specification

Assay min. 20%

For further information see separate document: "Standard Specification" (not for regulatory purposes) available via BASF's WorldAccount: <https://worldaccount.basf.com> (registered access).



$C_{40}H_{56}$

Molar mass 536.9 g/mol

Monographs

The active ingredient complies with the current "Beta-carotene" Ph. Eur. and "β-Carotene" FCC monographs, as well as the purity requirements of Directive 2008/128/EC (E160 a).

Regulations

Beta-carotene is approved for use as a food colorant and as a provitamin A source in most countries. However, specific regulations on the ingredients used in the respective countries and for the intended use have to be observed.

Bulk density

Approx. 0.6 g/ml

Stability

The product is stabilized with tocopherol (E306) and sodium ascorbate (E301).

The stability of Beta-Carotene 20% DC is excellent even in the presence of minerals. The product has a high mechanical integrity and little or none of the beta-carotene is expressed during tableting, resulting in good stability of the tablets. Stored in its original packaging at room temperature (max. 25 °C), it is stable for at least 36 months.

Storage/Handling

The product is sensitive to oxygen, light, heat, and moisture and should therefore be stored in the tightly closed original packaging at room temperature (max. 25 °C), in a dry and dark place.

Applications

Dietary supplements:

Beta-carotene dry powders are used as provitamin A in solid multivitamin preparations. Unlike vitamin A, no Upper Safe Level for use of beta-carotene as provitamin A has been established to date.

Beta-Carotene 20% DC is used both as a provitamin A and as an active ingredient. Due to its excellent flow properties, it is particularly suitable for use in multivitamin/mineral tablets and beta-carotene tablets, as well as hard gelatin capsules.

Food products:

Due to its composition, Beta-Carotene 20% DC is especially suitable for use in food supplement preparations, such as plain, effervescent, and sugar-coated tablets.

Note

Beta-Carotene 20% DC must be handled in accordance with the Safety Data Sheet.

This document, or any answers or information provided herein by BASF, does not constitute a legally binding obligation of BASF. While the descriptions, designs, data and information contained herein are presented in good faith and believed to be accurate, it is provided for your guidance only. Because many factors may affect processing or application/use, we recommend that you make tests to determine the suitability of a product for your particular purpose prior to use. It does not relieve our customers from the obligation to perform a full inspection of the products upon delivery or any other obligation.

NO WARRANTIES OF ANY KIND, EITHER EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE MADE REGARDING PRODUCTS DESCRIBED OR DESIGNS, DATA OR INFORMATION SET FORTH, OR THAT THE PRODUCTS, DESIGNS, DATA OR INFORMATION MAY BE USED WITHOUT INFRINGING THE INTELLECTUAL PROPERTY RIGHTS OF OTHERS. IN NO CASE SHALL THE DESCRIPTIONS, INFORMATION, DATA OR DESIGNS PROVIDED BE CONSIDERED A PART OF OUR TERMS AND CONDITIONS OF SALE.

November 2009